

Model with Strong γ_4 T -violation*

R. Friedberg¹ and T. D. Lee^{1, 2}

1. *Physics Department, Columbia University*

New York, NY 10027, U.S.A.

2. *China Center of Advanced Science and Technology (CCAST/World Lab.)*

P.O. Box 8730, Beijing 100080, China

Abstract

We extend the T violating model of the paper on "Hidden symmetry of the CKM and neutrino-mapping matrices" by assuming its T -violating phases $\chi_{\uparrow}$ and $\chi_{\downarrow}$ to be large and the same, with $\chi = \chi_{\uparrow} = \chi_{\downarrow}$. In this case, the model has 9 real parameters: $\alpha_{\uparrow}, \beta_{\uparrow}, \xi_{\uparrow}, \eta_{\uparrow}$ for the $\uparrow$ -quark sector, $\alpha_{\downarrow}, \beta_{\downarrow}, \xi_{\downarrow}, \eta_{\downarrow}$ for the $\downarrow$ sector and a common χ . We examine whether these nine parameters are compatible with ten observables: the six quark masses and the four real parameters that characterize the CKM matrix (i.e., the Jarlskog invariant $\mathcal{J}$ and three Eulerian angles). We find that this is possible only if the T -violating phase χ is large, between -120° to -135° . In this strong T violating model, the smallness of the Jarlskog invariant $\mathcal{J} \cong 3 \times 10^{-5}$ is mainly accounted for by the large heavy quark masses, with $\frac{m_c}{m_t} < \frac{m_s}{m_b} \approx .02$, as well as the near complete overlap of t and b quark, with $(c|b) = -.04$.

PACS: 12.15.Ff, 11.30.Er

Key words: Jarlskog invariant, CKM matrix, strong γ_4 T -violation

*This research was supported in part by the U.S. Department of Energy (grant no. DE-FG02-92-ER40699)

1. Introduction

In a previous paper on the "Hidden symmetry of the CKM and neutrino-mapping matrices" [1], we have posited a mass-generating Hamiltonian $H_\uparrow + H_\downarrow$ where

$$\begin{aligned} H_\uparrow &= \alpha_\uparrow |q_3^\uparrow - \xi_\uparrow q_2^\uparrow|^2 + \beta_\uparrow |q_2^\uparrow - \eta_\uparrow q_1^\uparrow|^2 + \beta_\uparrow |q_3^\uparrow - \xi_\uparrow \eta_\uparrow q_1^\uparrow|^2 \\ H_\downarrow &= \alpha_\downarrow |q_3^\downarrow - \xi_\downarrow q_2^\downarrow|^2 + \beta_\downarrow |q_2^\downarrow - \eta_\downarrow q_1^\downarrow|^2 + \beta_\downarrow |q_3^\downarrow - \xi_\downarrow \eta_\downarrow q_1^\downarrow|^2 \end{aligned} \quad (1.1)$$

with α, β, ξ, η real. This conserves T and leads to zero masses for the light quarks u and d . We then modified (1.1) by replacing $\xi_\uparrow, \xi_\downarrow$ with the corresponding T violating factors $\xi_\uparrow e^{i\chi_\uparrow}$ and $\xi_\downarrow e^{i\chi_\downarrow}$. To first order in $\chi_\uparrow$ and $\chi_\downarrow$ we found a relation of proportionality between $\mathcal{J}$, the Jarlskog invariant measuring T -violation, and a linear combination of square roots of the light masses. The ratio agreed roughly with known values. We shall call this the "weak γ_4 -model" because to make the calculation we assumed $\chi_\uparrow, \chi_\downarrow$ to be small.

There were two reasons for dissatisfaction with this model. First, why not introduce the phase factor into η or $\xi\eta$, yielding different physics? And second, when we estimated not only $\mathcal{J}$ but the individual matrix elements of U_{CKM} , we found that the data required $\chi_\uparrow$ and $\chi_\downarrow$ to be large angles, not small.

We now present a new model, the "strong γ_4 -model". In this model we introduce phase factors independently into all three terms, but require them to have the same values in $H_\uparrow$ and $H_\downarrow$. Thus we take the mass-generating Hamiltonian to be $H_\uparrow + H_\downarrow$ where

$$\begin{aligned} H_\uparrow &= \alpha_\uparrow |q_3^\uparrow - \xi_\uparrow e^{i\rho} q_2^\uparrow|^2 + \beta_\uparrow |q_2^\uparrow - \eta_\uparrow e^{i\omega} q_1^\uparrow|^2 + \beta_\uparrow |q_3^\uparrow - \xi_\uparrow \eta_\uparrow e^{-i\tau} q_1^\uparrow|^2 \\ H_\downarrow &= \alpha_\downarrow |q_3^\downarrow - \xi_\downarrow e^{i\rho} q_2^\downarrow|^2 + \beta_\downarrow |q_2^\downarrow - \eta_\downarrow e^{i\omega} q_1^\downarrow|^2 + \beta_\downarrow |q_3^\downarrow - \xi_\downarrow \eta_\downarrow e^{-i\tau} q_1^\downarrow|^2 \end{aligned} \quad (1.2)$$

It is now easily seen that the masses and CKM matrix depend on the phases only through the sum $\chi = \rho + \omega + \tau$. Accordingly, without loss of generality, we set $\rho = \omega = 0, \tau = \chi$. The mass-generating Hamiltonian can then be written as

$$\left(\bar{q}_1^\uparrow, \bar{q}_2^\uparrow, \bar{q}_3^\uparrow \right) M_\uparrow \begin{pmatrix} q_1^\uparrow \\ q_2^\uparrow \\ q_3^\uparrow \end{pmatrix} + \left(\bar{q}_1^\downarrow, \bar{q}_2^\downarrow, \bar{q}_3^\downarrow \right) M_\downarrow \begin{pmatrix} q_1^\downarrow \\ q_2^\downarrow \\ q_3^\downarrow \end{pmatrix}$$

where $q_i^\uparrow$, $q_i^\downarrow$ and $\bar{q}_i^\uparrow$, $\bar{q}_i^\downarrow$ are related to the corresponding Dirac field operators $\psi(q_i(\uparrow))$, $\psi(q_i(\downarrow))$ and their hermitian conjugate $\psi^\dagger(q_i(\uparrow))$, $\psi^\dagger(q_i(\downarrow))$ by

$$q_i^{\uparrow/\downarrow} = \psi(q_i(\uparrow / \downarrow)) \quad \text{and} \quad \bar{q}_i^{\uparrow/\downarrow} = \psi^\dagger(q_i(\uparrow / \downarrow))\gamma_4, \quad (1.3)$$

$$M_{\uparrow/\downarrow} = \begin{pmatrix} \beta\eta^2(1 + \xi^2) & -\beta\eta & -\beta\xi\eta e^{i\chi} \\ -\beta\eta & \beta + \alpha\xi^2 & -\alpha\xi \\ -\beta\xi\eta e^{-i\chi} & -\alpha\xi & \alpha + \beta \end{pmatrix}_{\uparrow/\downarrow}, \quad (1.4)$$

with the arrow-subscripts $\uparrow$, $\downarrow$ referring to α , β , ξ , η , but not to χ .

In diagonalizing (1.4) we do not assume, as in the weak γ_4 -model, that χ is small. We find that the smallness of $\mathcal{J}$ is mainly accounted for by the large heavy masses with

$$\frac{m_c}{m_t} < \frac{m_s}{m_b} \approx .02 \quad (1.5)$$

and by the nearly complete overlap of the statevectors for t and b since

$$|(u|b)| < |(c|b)| \cong 0.04. \quad (1.6)$$

We have been able to carry out complete calculations in which the only approximations are based on the smallness of $\frac{m_s}{m_b}$, $\frac{m_c}{m_t}$ and $(c|b)$. These calculations are described in Sections 2 and 3; we give here a brief outline.

We diagonalize $M_\uparrow$ and $M_\downarrow$ with the aid of parameters $r_{\uparrow,\downarrow}$, $B_{\uparrow,\downarrow}$, $\Phi_{\uparrow,\downarrow}$, $\mathcal{S}$, $\mathcal{L}$ to be defined in the next two sections. They are shown there to satisfy the following ten equations (to first order in small quantities):

$$\frac{1 - r_\uparrow^2}{r_\uparrow^2} \sin^2 B_\uparrow = \frac{4m_u m_c}{(m_c - m_u)^2}, \quad (1.7)$$

$$\frac{1 - r_\downarrow^2}{r_\downarrow^2} \sin^2 B_\downarrow = \frac{4m_d m_s}{(m_s - m_d)^2}, \quad (1.8)$$

$$\sin^2 \frac{1}{2}\chi = \frac{1 - r_\uparrow^2}{\sin^2 2\Phi_\uparrow} = \frac{1 - r_\downarrow^2}{\sin^2 2\Phi_\downarrow}, \quad (1.9)$$

$$\mathcal{L} = \frac{\sqrt{m_s m_d}}{m_b} - \frac{\sqrt{m_c m_u}}{m_t}, \quad (1.10)$$

$$\mathcal{S} = \sin(\Phi_\uparrow - \Phi_\downarrow) = (c|b), \quad (1.11)$$

$$|(u|b) + \mathcal{S} \sin \frac{1}{2} B_{\uparrow}|^2 = \mathcal{L}^2 \cos^2 \frac{1}{2} B_{\uparrow}, \quad (1.12)$$

$$Im(u|b) = -\mathcal{L} \frac{\cos \frac{1}{2} B_{\uparrow} \cos \frac{1}{2} \chi}{r_{\uparrow}} \quad (1.13)$$

and

$$(u|s) = \sin \frac{1}{2} (B_{\downarrow} - B_{\uparrow}). \quad (1.14)$$

Our strategy of solution is as follows. We take m_s , m_c , m_b , m_t , as well as $(u|s)$, $(u|b)$ and $(c|b)$, to be given from data (see table 1). Then we have eleven unknowns $r_{\uparrow, \downarrow}$, $B_{\uparrow, \downarrow}$, $\Phi_{\uparrow, \downarrow}$, $\mathcal{S}$, $\mathcal{L}$, χ , m_d , m_u constrained by ten independent equations given above (with (1.9) and (1.11), each counted as two equations). Taking a trial value of $\sin \frac{1}{2} B_{\uparrow}$, we are able to solve numerically for the other ten unknowns by a self-correcting double iteration that converges to 4 decimal stability after $36 = 6 \times 6$ passes. We find that m_u is particularly sensitive to variations in $\sin \frac{1}{2} B_{\uparrow}$; a variation of 30% in the latter carries m_u through the whole of its experimental range from 1.5 to $3.0 MeV/c^2$. Meanwhile m_d varies by only 25%, from 5.2 to $6.5 MeV/c^2$, well within the experimental range, 3.0 to $8.0 MeV/c^2$. The value of χ must be taken as negative and is in the neighborhood of -125° , between -120° and -135° . We have also tried deviations in m_s , m_b , $(c|b)$, $Re(u|b)$ and $Im(u|b)$. Only in the case of m_s does it appear that a maximal deviation (-25%) from the "best value" might push m_d outside the range given by data. (See Tables 1 and 2, and Fig. 1).

The next two sections are devoted to defining the parameters that appear in (1.7)-(1.14) and proving that these equations are satisfied. In Section 2, we discuss the separate diagonalization of $M_{\uparrow}$ and $M_{\downarrow}$, and in Section 3, we examine the CKM matrix.

In Section 4, we discuss briefly a third model[2], which we may call a $i\gamma_5$ model, because its Hamiltonian contains a term in $i\gamma_4\gamma_5$ as well as the usual one in γ_4 .

2. Diagonalization of $M_\uparrow$ and $M_\downarrow$

In this section, we shall drop the arrow-subscripts and write (1.4) as

$$M = \begin{pmatrix} T^2\beta & -T\beta \cos \Phi & -T\beta \sin \Phi e^{i\chi} \\ -T\beta \cos \Phi & \alpha \tan^2 \Phi + \beta & -\alpha \tan \Phi \\ -T\beta \sin \Phi e^{-i\chi} & -\alpha \tan \Phi & \alpha + \beta \end{pmatrix}, \quad (2.1)$$

where

$$\Phi = \tan^{-1} \xi \quad (2.2)$$

$$T = \eta \sqrt{1 + \xi^2} \quad (2.3)$$

so that $T^2\beta = \beta\eta^2(1 + \xi^2)$, $\sin \Phi = \xi/\sqrt{1 + \xi^2}$, $\cos \Phi = 1/\sqrt{1 + \xi^2}$ and (2.1)=(1.4). We denote the eigenvalues of M by m_l , m_m , m_h (light, medium, heavy), and seek a unitary matrix W (with $WW^\dagger = 1$) such that

$$M = W \begin{pmatrix} m_l & 0 & 0 \\ 0 & m_m & 0 \\ 0 & 0 & m_h \end{pmatrix} W^\dagger. \quad (2.4)$$

The W matrix will be built up in stages, as we shall discuss. First we isolate the heavy mass by writing

$$M = \Omega \left(\begin{array}{c|c} (n) & L \\ \hline L^* & 0 \end{array} \right) \Omega^\dagger \quad (2.5)$$

where

$$\Omega^\dagger = \left(\begin{array}{c|cc} 1 & 0 & 0 \\ \hline 0 & e^{i\Phi\tau_y} & \\ 0 & & \end{array} \right), \quad (2.6)$$

$$\mu_h = \alpha \sec^2 \Phi + \beta \quad (2.7)$$

$$L = T\beta \cos \Phi \sin \Phi (1 - e^{i\chi}) \quad (2.8)$$

and

$$(n) = \beta \begin{pmatrix} T^2 & -T(\cos^2 \Phi + \sin^2 \Phi e^{i\chi}) \\ -T(\cos^2 \Phi + \sin^2 \Phi e^{-i\chi}) & 1 \end{pmatrix}. \quad (2.9)$$

Thus, (2.1) can be obtained by a simple substitution of (2.6)-(2.9) into (2.5).

Next, we diagonalize the 2×2 matrix (n) of (2.9) by setting

$$\cos^2 \Phi + \sin^2 \Phi e^{i\chi} = r e^{iA} \quad (2.10)$$

with r, A both real. Then

$$\begin{aligned} (n) &= \beta \begin{pmatrix} T^2 & -T r e^{iA} \\ -T r e^{-iA} & 1 \end{pmatrix} \\ &= e^{\frac{1}{2}i\tau_z A} e^{-\frac{1}{2}i\tau_y B} \begin{pmatrix} \mu_l & 0 \\ 0 & \mu_m \end{pmatrix} e^{\frac{1}{2}i\tau_y B} e^{-\frac{1}{2}i\tau_z A} \end{aligned} \quad (2.11)$$

provided that

$$\begin{aligned} \mu_m + \mu_l &= \beta(1 + T^2) \\ (\mu_m - \mu_l) \cos B &= \beta(1 - T^2) \\ (\mu_m - \mu_l) \sin B &= 2\beta T r. \end{aligned} \quad (2.12)$$

By quadratic combination of (2.12) we obtain

$$\mu_m \mu_l = \beta^2 T^2 (1 - r^2); \quad (2.13)$$

then, by dividing the above equation by the square of the last line of (2.12), we have

$$\frac{4\mu_m \mu_l}{(\mu_m - \mu_l)^2} = \frac{1 - r^2}{r^2} \sin^2 B \quad (2.14)$$

which leads to (1.7) and (1.8).

Also, by applying the Law of Sines to the complex triangle described by (2.10), followed by trigonometric identities, we find

$$\cos\left(\frac{1}{2}\chi - A\right) = \frac{\cos \frac{1}{2}\chi}{r}, \quad (2.15)$$

a relation that will be useful later.

Applying (2.11) to (2.5), we now have

$$M = \Omega V \begin{pmatrix} \mu_l & 0 & L\Delta^* \cos \frac{1}{2}B \\ 0 & \mu_m & -L\Delta^* \sin \frac{1}{2}B \\ L^* \Delta \cos \frac{1}{2}B & -L^* \Delta \sin \frac{1}{2}B & \mu_h \end{pmatrix} V^\dagger \Omega^\dagger, \quad (2.16)$$

where

$$\Delta = e^{\frac{1}{2}iA} \quad (2.17)$$

and

$$V^\dagger = \left(\begin{array}{cc|c} \left(e^{\frac{1}{2}i\tau_y B} e^{-\frac{1}{2}i\tau_z A} \right) & & 0 \\ & & 0 \\ \hline 0 & 0 & 1 \end{array} \right) \quad (2.18)$$

Thus M is almost diagonalized. Let us study the magnitude of L . From (2.13) and (2.10) we find

$$\mu_m \mu_l = \beta^2 T^2 (1 - r^2) = 2\beta^2 T^2 (1 - \cos \chi) \cos^2 \Phi \sin^2 \Phi \quad (2.19)$$

and comparing this with (2.8) we have

$$|L| = 2|T\beta \cos \Phi \sin \Phi \sin \frac{1}{2}\chi| = \sqrt{\mu_m \mu_l}. \quad (2.20)$$

Hence, if we write

$$\begin{pmatrix} \mu_l & 0 & L\Delta^* \cos \frac{1}{2}B \\ 0 & \mu_m & -L\Delta^* \sin \frac{1}{2}B \\ L^* \Delta \cos \frac{1}{2}B & -L^* \Delta \sin \frac{1}{2}B & \mu_h \end{pmatrix} = P \begin{pmatrix} m_l & 0 & 0 \\ 0 & m_m & 0 \\ 0 & 0 & m_h \end{pmatrix} P^\dagger \quad (2.21)$$

the elements of P will differ from those of the unit matrix by $O\left[\frac{\sqrt{m_l m_m}}{m_h}\right] \ll 1$. A careful examination shows that all the m 's may be approximated by μ 's; in particular, we also have $|\frac{\mu_l}{m_l} - 1| \sim O\left[\frac{m_m}{m_h}\right]$. Therefore (2.14) becomes

$$\frac{4m_m m_l}{(m_m - m_l)^2} = \frac{1 - r^2}{r^2} \sin^2 B \quad (2.22)$$

and (1.7) and (1.8) are established.

Also, (1.9) is a direct consequence of (2.13) and (2.20). We may take (1.10) as the definition of $\mathcal{L}$, and from (2.20) we may write it as

$$\mathcal{L} = \frac{|L_\downarrow|}{m_b} - \frac{|L_\uparrow|}{m_t}. \quad (2.23)$$

The first equality of (1.11) is the definition of $\mathcal{S}$. Thus what remains is to establish the second part of (1.11), and (1.12)-(1.14). This requires studying the CKM matrix which relates "up" to "down" eigenstates, as we shall see.

3. The CKM Matrix

In this section we restore the arrow subscripts $\uparrow, \downarrow$. On account of (2.16) and (2.21), the matrix W defined in (2.4) is given by

$$W_{\uparrow,\downarrow}^\dagger = P_{\uparrow,\downarrow}^\dagger V_{\uparrow,\downarrow}^\dagger \Omega_{\uparrow,\downarrow}^\dagger. \quad (3.1)$$

If we define

$$U = W_\uparrow^\dagger W_\downarrow = P_\uparrow^\dagger U_0 P_\downarrow \quad (3.2)$$

where

$$\begin{aligned} U_0 &= V_\uparrow^\dagger \Omega_\uparrow^\dagger \Omega_\downarrow V_\downarrow \\ &= \left(\begin{array}{cc|c} \left(e^{\frac{1}{2}i\tau_y B_\uparrow} e^{-\frac{1}{2}i\tau_z A_\uparrow} \right) & 0 & \\ \hline 0 & 0 & 1 \end{array} \right) \left(\begin{array}{cc|c} 1 & 0 & 0 \\ \hline 0 & e^{i(\Phi_\uparrow - \Phi_\downarrow)\tau_y} & 0 \\ \hline 0 & 0 & 1 \end{array} \right) \left(\begin{array}{cc|c} \left(e^{\frac{1}{2}i\tau_z A_\downarrow} e^{-\frac{1}{2}i\tau_y B_\downarrow} \right) & 0 & \\ \hline 0 & 0 & 1 \end{array} \right) \end{aligned} \quad (3.3)$$

then U transforms eigenstates of $M_\downarrow$ into eigenstates of $M_\uparrow$, provided that the phases of the eigenstates are suitably chosen. To obtain the CKM matrix U_{CKM} , which relates eigenstates whose phases follow a standard convention, we shall need an additional transformation

$$U_{CKM} = Q_\uparrow^\dagger U Q_\downarrow \quad (3.4)$$

where $Q_{\uparrow,\downarrow}$ are diagonal unitary matrices to be chosen presently.

In evaluating (3.3) it is convenient to introduce new symbols:

$$\delta = \Delta_\uparrow \Delta_\downarrow^* = e^{\frac{1}{2}i(A_\uparrow - A_\downarrow)}, \quad (3.5)$$

$$\Gamma = \cos \frac{1}{2} B_\uparrow, \quad \gamma = \cos \frac{1}{2} B_\downarrow, \quad (3.6)$$

$$\Sigma = \sin \frac{1}{2} B_\uparrow, \quad \sigma = \sin \frac{1}{2} B_\downarrow, \quad (3.7)$$

$$\mathcal{S} = \sin(\Phi_\uparrow - \Phi_\downarrow) \text{ and } C = \cos(\Phi_\uparrow - \Phi_\downarrow). \quad (3.8)$$

We note that the first equation in (3.8) is the same in (1.11). By using (3.5)-(3.8), we find U_0 of (3.3) can be written as

$$U_0 = \begin{pmatrix} \delta^* \Gamma \gamma + C \delta \Sigma \sigma & -\delta^* \Gamma \sigma + C \delta \Sigma \gamma & \mathcal{S} \Delta_\uparrow \Sigma \\ -\delta^* \Sigma \gamma + C \delta \Gamma \sigma & \delta^* \Sigma \sigma + C \delta \Gamma \gamma & \mathcal{S} \Delta_\uparrow \Gamma \\ -\mathcal{S} \Delta_\downarrow^* \sigma & -\mathcal{S} \Delta_\downarrow^* \gamma & C \end{pmatrix}. \quad (3.9)$$

The next step is to prepare for a perturbative treatment of (3.2) by writing

$$P_{\uparrow,\downarrow} \cong I + p_{\uparrow,\downarrow} \quad (3.10)$$

where (in arrowless notation)

$$p^\dagger = \frac{1}{m_h} \begin{pmatrix} 0 & 0 & -\Delta^* L \cos \frac{1}{2} B \\ 0 & 0 & \Delta^* L \sin \frac{1}{2} B \\ \Delta L^* \cos \frac{1}{2} B & -\Delta L^* \sin \frac{1}{2} B & 0 \end{pmatrix}. \quad (3.11)$$

We note that by putting (3.11) into (3.10), we can satisfy (2.21) to first order in L .

Thus we have

$$U \cong U_0 + U' \quad (3.12)$$

where

$$U' = p^\dagger U_0 + U_0 p_\downarrow. \quad (3.13)$$

Let us carefully evaluate the lower left element of $p^\dagger U_0$:

$$\begin{aligned} (p^\dagger U_0)_{31} &= \frac{1}{m_t} (L_\uparrow^* \Delta_\uparrow \cos \frac{1}{2} B_\uparrow) (\delta^* \Gamma \gamma + C \delta \Sigma \sigma) \\ &\quad + \frac{1}{m_t} (-L_\uparrow^* \Delta_\uparrow \sin \frac{1}{2} B_\uparrow) (-\delta^* \Sigma \gamma + C \delta \Gamma \sigma) \\ &= \frac{L_\uparrow^* \Delta_\uparrow}{m_t} [\Gamma (\delta^* \Gamma \gamma + C \delta \Sigma \sigma) + \Sigma (\delta^* \Sigma \gamma - C \delta \Gamma \sigma)] \\ &= \frac{L_\uparrow^* \Delta_\uparrow}{m_t} \delta^* (\Gamma^2 + \Sigma^2) \gamma = \frac{L_\uparrow^*}{m_t} \Delta_\downarrow \gamma. \end{aligned} \quad (3.14)$$

(Note how the calculation converts $\Delta_\uparrow$ to $\Delta_\downarrow$ and Γ to γ .) The corresponding element of $U_0 p_\downarrow$ is trivial:

$$(U_0 p_\downarrow)_{31} = C \left(\frac{1}{m_b} \Delta_\downarrow^* L_\downarrow \cos \frac{1}{2} B_\downarrow \right)^* = -\frac{L_\downarrow^*}{m_b} \Delta_\downarrow \gamma C. \quad (3.15)$$

Anticipating that $B_\uparrow$ will turn out fairly small, ~ 0.2 , we now observe that the matrix element U_{23} is going to be dominated by $(U_0)_{23} = \mathcal{S} \Delta_\uparrow \Gamma \sim \mathcal{S} \Delta_\uparrow$. Therefore, $\mathcal{S}$ must have magnitude $\sim .04$. It follows that $C \sim 1 - \frac{1}{2} \mathcal{S}^2$ can

be replaced by 1, and that all elements of U' other than $(U')_{13,23,31,32}$ being of order $\mathcal{S} \cdot \frac{\sqrt{m_d m_s}}{m_b}$, can be neglected.

Thus, by repeating for $(U')_{13,23,32}$ the calculations leading to (3.14) and (3.15), we have

$$U' \cong \begin{pmatrix} 0 & 0 & +(\frac{L_\downarrow}{m_b} - \frac{L_\uparrow}{m_t})\Delta_\uparrow\Gamma \\ 0 & 0 & -(\frac{L_\downarrow}{m_b} - \frac{L_\uparrow}{m_t})\Delta_\uparrow\Sigma \\ -(\frac{L_\downarrow^*}{m_b} - \frac{L_\uparrow^*}{m_t})\Delta_\downarrow\gamma & +(\frac{L_\downarrow^*}{m_b} - \frac{L_\uparrow^*}{m_t})\Delta_\downarrow\sigma & 0 \end{pmatrix}. \quad (3.16)$$

But from (2.8), taking $T, \beta, \cos \Phi, \sin \Phi$ positive, we find

$$\frac{L_\downarrow}{|L_\downarrow|} = \frac{L_\uparrow}{|L_\uparrow|} = \frac{1 - e^{i\chi}}{|1 - e^{i\chi}|} \quad (3.17)$$

and so

$$\frac{L_\downarrow}{m_b} - \frac{L_\uparrow}{m_t} = \frac{1 - e^{i\chi}}{|1 - e^{i\chi}|} \mathcal{L} \quad (3.18)$$

by (2.23). We now anticipate that χ will have to be negative in order to make everything come out right. Hence,

$$\frac{1 - e^{i\chi}}{|1 - e^{i\chi}|} = \frac{e^{\frac{1}{2}i\chi}(-2i \sin \frac{1}{2}\chi)}{|2 \sin \frac{1}{2}\chi|} = +ie^{\frac{1}{2}i\chi} \quad (3.19)$$

and (3.16) leads to

$$U' \simeq \begin{pmatrix} 0 & 0 & +ie^{\frac{1}{2}i\chi}\mathcal{L}\Delta_\downarrow^*\Gamma \\ 0 & 0 & -ie^{\frac{1}{2}i\chi}\mathcal{L}\Delta_\downarrow^*\Sigma \\ +ie^{-\frac{1}{2}i\chi}\mathcal{L}\Delta_\downarrow\gamma & -ie^{-\frac{1}{2}i\chi}\mathcal{L}\Delta_\downarrow\sigma & 0 \end{pmatrix}. \quad (3.20)$$

For reasons shortly to be evident, let us now introduce the phase factors

$$\varepsilon_{\uparrow,\downarrow} = -ie^{\frac{1}{2}i\chi}(\Delta_{\uparrow,\downarrow}^*)^2 = e^{-\frac{i\sigma}{2}}e^{i(\frac{1}{2}\chi - A_{\uparrow,\downarrow})}. \quad (3.21)$$

Then we have

$$U' = \begin{pmatrix} 0 & 0 & -\varepsilon_\uparrow\mathcal{L}\Delta_\uparrow\Gamma \\ 0 & 0 & +\varepsilon_\uparrow\mathcal{L}\Delta_\uparrow\Sigma \\ +\varepsilon_\downarrow^*\mathcal{L}\Delta_\downarrow^*\gamma & -\varepsilon_\downarrow^*\mathcal{L}\Delta_\downarrow^*\sigma & 0 \end{pmatrix}. \quad (3.22)$$

In treating (3.9), let us note that since $\Phi_\uparrow - \Phi_\downarrow \approx \sin^{-1} \mathcal{S}$ is small, $A_\uparrow - A_\downarrow$ is also small by (2.10). Hence $|Im \delta|$ is small (see(3.5)) and $1 - Re \delta$ is second order. So $Re \delta$ can be taken = 1, and the imaginary parts of $(U_0)_{11,12,21,22}$ can be adjusted by small adjustments in $Q_\uparrow, Q_\downarrow$. We shall treat such adjustments imprecisely and simply neglect these imaginary parts. By taking $C \rightarrow 1$ and using (3.6)-(3.7), we find

$$\begin{pmatrix} (U_0)_{11} & (U_0)_{12} \\ (U_0)_{21} & (U_0)_{22} \end{pmatrix} = \begin{pmatrix} \Gamma\gamma + \Sigma\sigma & -\Gamma\sigma + \Sigma\gamma \\ -\Sigma\gamma + \Gamma\sigma & \Sigma\sigma + \Gamma\gamma \end{pmatrix} \\ = \begin{pmatrix} \cos \frac{1}{2}(B_\downarrow - B_\uparrow) & -\sin \frac{1}{2}(B_\downarrow - B_\uparrow) \\ \sin \frac{1}{2}(B_\downarrow - B_\uparrow) & \cos \frac{1}{2}(B_\downarrow - B_\uparrow) \end{pmatrix}. \quad (3.23)$$

Now $B_\downarrow - B_\uparrow$ must be positive to fit U_{13} and U_{31} , and so U_{12} is negative, whereas the standard presentation gives $(U_{CKM})_{12}$ positive. Therefore, we shall use the Q -transformation to change the sign of the first row and column, and also to remove the factors $\Delta_\uparrow, \Delta_\downarrow^*$ now appearing in the third row and column. Thus

$$Q_\uparrow^\dagger = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \Delta_\downarrow \end{pmatrix}, \quad Q_\downarrow^\dagger = \begin{pmatrix} +1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \Delta_\uparrow^* \end{pmatrix} \quad (3.24)$$

and

$$U_{CKM} = Q_\uparrow^\dagger U_0 Q_\downarrow + Q_\uparrow^\dagger U' Q_\downarrow \\ = \begin{pmatrix} \cos \frac{1}{2}(B_\downarrow - B_\uparrow) & \sin \frac{1}{2}(B_\downarrow - B_\uparrow) & -\mathcal{S}\Sigma + \varepsilon_\uparrow \mathcal{L}\Gamma \\ -\sin \frac{1}{2}(B_\downarrow - B_\uparrow) & \cos \frac{1}{2}(B_\downarrow - B_\uparrow) & \mathcal{S}\Gamma + \varepsilon_\uparrow \mathcal{L}\Sigma \\ \mathcal{S}\sigma - \varepsilon_\downarrow^* \mathcal{L}\gamma & -\mathcal{S}\gamma - \varepsilon_\downarrow^* \mathcal{L}\sigma & 1 \end{pmatrix} \quad (3.25)$$

where we have again allowed a slight imprecision of phase in the (3, 3) element.

Comparing (3.25) with the array

$$U_{CKM} = \begin{pmatrix} (u|d) & (u|s) & (u|b) \\ (c|d) & (c|s) & (c|b) \\ (t|d) & (t|s) & (t|b) \end{pmatrix}, \quad (3.26)$$

we obtain the second half of (1.11) and (1.12)-(1.14).

Note: there is an ambiguity, $\Phi_{\uparrow,\downarrow} > \text{ or } < \frac{\pi}{4}$. We take both Φ 's $> \frac{\pi}{4}$, so that $|A| > |\chi - A|$ and hence $|A| > |\frac{1}{2}\chi|$. Since χ and A are negative, $\frac{1}{2}\chi - A > 0$ and hence $Re \varepsilon_{\uparrow,\downarrow} > 0$, as required in $(u|b)$ and $(t|d)$. Because $Im \varepsilon_{\uparrow} = -\cos(\frac{1}{2}\chi - A)$, we can then derive (1.13) by using (2.15).

4. The "Timeon" Model

The merit of the "strong γ_4 T -violation model" examined in this paper suggests that there may be large T -violation somewhere in physics although its manifestation in the quark mass sector is small. In the "strong γ_4 T -violation model" the T -violating effects are produced by the phase χ which enters non-linearly into the Hamiltonian. This non-linear interaction makes it difficult to construct a renormalizable quantum field theory that can be extended beyond the mass matrix. For this and other reasons, we have considered a different model[3] in which the T -violating effect enters linearly; therefore, the model can lead to a renormalizable field theory, called "timeon".

In the timeon theory, the mass-generating Hamiltonian can be written by replacing $M_{\uparrow/\downarrow}$ in (1.4) by

$$G_{\uparrow/\downarrow} + i\gamma_5 F_{\uparrow/\downarrow}, \quad (4.1)$$

where $G_{\uparrow/\downarrow}$ and $F_{\uparrow/\downarrow}$ are real symmetric matrices, and the $F_{\uparrow/\downarrow}$ term in $i\gamma_4$ arises from coupling to the vacuum expectation value of a new T -negative and P -negative field $\tau(x)$, the timeon field. Thus, the whole field theory conserves T , but T -violation arises from the spontaneous symmetry breaking that makes the vacuum expectation value

$$\tau_0 = \langle \tau(x) \rangle_{vac} \neq 0. \quad (4.2)$$

The timeon field $\tau(x)$ is real, so that there is no Goldstone boson[4]. However, the oscillation of $\tau(x)$ around its vacuum expectation value τ_0 gives rise to a new particle, called "timeon", whose production can lead to large T -violating effects. In Ref. 3, it is shown that the parameters determining $G_{\uparrow/\downarrow}$ and $F_{\uparrow/\downarrow}$ can be adjusted to simulate an arbitrary complex γ_4 model, as far as the quark masses are concerned, but not the CKM matrix. Thus, for example,

in the timeon γ_5 -model the light quark masses in the small mass limit turn out to be proportional to $\mathcal{J}$, whereas in the γ_4 -model, they are proportional to $\mathcal{J}^2$.

References

- [1] R. Friedberg and T. D. Lee, Ann. Phys. **323**(2008)1087
- [2] Particle Data Group, J. Phys. **G33**(2006)1
- [3] R. Friedberg and T. D. Lee, arXiv:0809.3633
- [4] J. Goldstone, Nuovo Cimento **9**(1961)154

Table 1

Parameter	"Best" value
m_s	$95MeV$
m_b	$4.5GeV$
$(c b)$	0.04
$Re(u b)$	0.002
$Im(u b)$	-0.003

These values are used to obtain the top two rows in Table 2.

Table 2

Values of m_u , m_d and χ calculated from the strong γ_4 -model

Input parameters		$m_u(MeV)$	$m_d(MeV)$	$\cos \frac{1}{2}\chi$
As in Table 1		1.45	5.18	.487
		3.16	6.50	.428
Table 1 except	$m_s = 85MeV$	1.39	5.43	.479
		3.29	6.86	.418
Table 1 except	$m_s = 105MeV$	1.52	5.00	.490
		3.09	6.22	.433
Table 1 except	$m_b = 4.2GeV$	1.63	4.83	.483
		3.33	6.02	.427
Table 1 except	$m_b = 4.7GeV$	1.61	5.68	.476
		3.53	7.14	.417
Table 1 except	$(c b) = 0.035$	1.40	4.86	.507
		2.98	5.96	.454
Table 1 except	$(c b) = 0.045$	1.51	5.52	.468
		3.36	7.07	.405
Table 1 except	$Re(u b) = 0.0015$	1.63	4.74	.525
		3.33	5.96	.463
Table 1 except	$Re(u b) = 0.0025$	1.72	6.09	.432
		2.96	7.06	.397
Table 1 except	$Im(u b) = -0.0025$	1.64	4.93	.428
		2.75	5.81	.389
Table 1 except	$Im(u b) = -0.0035$	1.73	5.96	.510
		2.93	6.83	.473

Table 2 (footnotes)

The values of five input parameters are taken as in Table 1, except for single departures as shown in the left-hand column here. For each setting of the input parameters, there is a one-parameter family of solutions of Eqs. (1.7)-(1.14). We show two members of each family, chosen roughly to span the experimental range of m_u from 1.5 to 3.0 MeV . The corresponding values of m_d stay within its experimental range from 3 to 8 MeV , and χ remains large from -120° to -135° .




Figure 1. m_d versus m_u for $m_s = 95\text{MeV}$, $(c|b) = 0.04$,
 $(u|b) = 0.002 - 0.003i$ and $m_b = 4.2\text{GeV}$, 4.5GeV and 4.7GeV .